

Public Health Services for Berkshire

Berkshire Suicide Prevention Strategy 2017-2020

Full Version with Audit and Action Plans

DRAFT V10


Darrell Gale FFPH

Consultant in Public Health

Mental Health Lead Consultant for Berkshire



WOKINGHAM
BOROUGH COUNCIL



DIGITAL SOLUTIONS PROOF

Contents

Acknowledgments	4
Executive Summary	5
Recommendations	6
Background	8
10 Things Everyone Needs To Know About Suicide Prevention	9
Strategy Aims	10
National Context	11
Strategic Context	14
Evidence Base in Suicide Prevention	15
National Best Practice in Suicide Prevention.....	16
Local Context	18
Local Suicide Audit Results	18
Local Governance Structures	28
Local Best Practice in Suicide Prevention	30
Areas of High Frequency.....	32
Crisis Care Concordat	35
Gap Analysis and Emergent Berkshire-Wide Concerns	36
Berkshire-Wide Action Plan 2017-18	39
References.....	43

Appendices

Appendix 1 Resources available.....	44
Appendix 2 Bracknell Forest Action Plan 2017-18.....	45
Appendix 3 RBWM Action Plan 2017-18	47
Appendix 4 Slough Action Plan 2017-18.....	50
Appendix 5 Reading Action Plan 2017-18	53
Appendix 6 West Berkshire Action Plan 2017-18.....	56
Appendix 7 Wokingham Action Plan 2017-18.....	58
Appendix 8 Membership of the Berkshire Suicide Prevention Steering Group	61

Acknowledgements

Acknowledgements are due to a wide range of partners and colleagues whose work; encouragement and commitment to suicide prevention has enabled the development of this strategy and its action plans. In particular, we acknowledge the following:

Rutuja Kulkarni and the public health officers from Berkshire local authorities who undertook the Suicide Audit, and who did much to build the foundations upon which this strategy has developed;

The suicide prevention and mental health leads from the Berkshire local authorities for preparing the local action plans;

Network Rail and British Transport Police for their support with work on railway suicides;

Helena Fahie at Public Health England South East Centre for encouragement; advice and going the extra mile;

David Colchester at the Local Criminal Justice Board for the Thames Valley and Thames Valley Police for their input on real-time surveillance;

The NHS Provider trusts in Berkshire for their input and continued support;

The seven Clinical Commissioning Groups in Berkshire for their strong partnership working;

And of course all past and present members of the Berkshire Suicide Prevention Steering Group.

DIGITAL
SOLUTIONS
PROOF

Executive Summary

The NHS Five Year Forward View for Mental Health sets a target on all NHS agencies and partners to reduce the current level of suicide by 10% by 2020. This is a laudable and hopefully readily achievable aim. However as discussions across the range of organisations which have contributed to this strategy have progressed, it appears to many, that this aim is not challenging enough. Zero Suicide should be our aim; as it is in the gift of the combined efforts of these organisations, and of society at large, to put in place the policies and services which protect people from mental distress, and to ease the factors which cause that distress. This strategy therefore forges ahead with an ambition to reduce suicide by at least 25% by 2020, thus ensuring that this becomes a shared priority across organisations and areas.

We recognise that a Berkshire without suicide is the true aim we work towards.

This strategy is an important public health strategy which seeks to save lives lost to suicide through its prevention, and to improve the health and wellbeing of those bereaved by suicide. It also includes more general whole-population actions aimed at improving mental health and wellbeing as contributing factors that prevent suicide. The strategy highlights, and action plans prioritise, certain population groups which have greater risk factors for suicide, and thus contributes to narrowing health inequalities.

It goes without saying, but we should remind ourselves, that suicides are tragedies for all involved. For every person who dies by suicide at least 10 people are directly affected. Support for those bereaved, including the professionals who deal with the suicide, is vitally important. The social and economic cost of a suicide is substantial. The average cost of suicide in someone of working age in England is estimated to be £1.67 million. This includes direct costs of care, indirect costs relating to loss of productivity and earnings, and the intangible costs associated with pain, grief and suffering of those bereaved or affected by suicide.

Many stakeholders have contributed to this strategy and it should now be adopted as a joint strategy by each of the CCGs, Local Authorities, and the Health and Wellbeing Boards in Berkshire. It should also be referenced and reflected in other plans and strategies when they are drafted or re-written, to ensure suicide prevention becomes a pursuit common to all agencies and professions. It is an important and happy coincidence that this strategy will be formally launched, once it has been endorsed by all health and wellbeing boards in Berkshire, during Brighter Berkshire, the Year of Mental Health. This community led initiative aims to help increase the opportunities and support for our Berkshire population who need help with their mental health, when they need it and to build a stronger happier Berkshire population. The aims of this strategy fit well with these broader aims.

Dr Lise Llewellyn
Strategic Director of Public Health for Berkshire April 2017

RECOMMENDATION

Launch this strategy at a multi-agency suicide prevention summit, by October 2017.

Recommendations

The following recommendations are the principle strategic objectives for Berkshire as a whole. These link through into more detailed action plans for Berkshire-wide work and for local authority areas. In line with the national suicide prevention strategy, the ambition of this strategy is to reduce suicides in Berkshire by 25% by 2020, and to provide better support for those bereaved or affected by suicide. The national strategy has identified six priority areas and the recommendations linked to these are outlined below, following those relating to the overarching aims.

Over-arching Recommendations

RECOMMENDATION

That this Steering Group revisit their terms of reference and membership and become known as the "Berkshire Suicide Prevention Steering Group", with the aim of providing the governance to this strategy and its action plans.

RECOMMENDATION

That organisations represented on the Berkshire Suicide Prevention Steering Group consider nominating a suicide prevention champion from within their membership.

RECOMMENDATION

That all agencies represented on the Steering Group commit to an annual Action Plan of their own which can then be brought together to create a Berkshire-wide Action Plan for each year of the Strategy.

RECOMMENDATION

Sign off / endorsement of this strategy by all Health & Wellbeing Boards in Berkshire.

RECOMMENDATION

Launch this strategy at a multi-agency suicide prevention summit, by October 2017.

RECOMMENDATION

Ensure strategic fit with the action plans of the Berkshire Crisis Care Concordat.

Priority Areas

1. Reduce the risk of suicide in key high-risk groups;

RECOMMENDATION

Implement the NICE guidelines on self-harm, specifically ensuring that people who present to Emergency Departments following self-harm receive a psychosocial assessment.

RECOMMENDATION

Evaluate the Berkshire CALMzone and re-commission targeted suicide prevention work for younger men and middle aged men.

2. Tailor approaches to improve mental health in specific groups;

RECOMMENDATION

Work to provide and commission interventions which improve the public's mental health. These may include: awareness of mental health and peer support in young people; anti-bullying campaigns in schools; addressing stigma and social isolation in older people; workplace health promotion and support with local business; working with police on mental health literacy; and addressing issues relevant to the local population.

3. Reduce access to the means of suicide;

RECOMMENDATION

That local authority public health teams take the leadership for liaison with any Escalation Process in their area, and report on progress to the Steering Group.

RECOMMENDATION

That the Steering Group seek a named Highways England officer to act as a liaison link and group member, and to share real-time intelligence of highways network incidents.

RECOMMENDATION

That local authority public health teams work with other council departments such as car parks and open space services to identify local actions to prevent suicide including staff awareness training.

4. Provide better information and support to those bereaved or affected by suicide;

RECOMMENDATION

Ensure bereavement information and access to support is available to those bereaved by suicide, including professionals involved in the case.

5. Support the media in delivering sensitive approaches to suicide and suicidal behaviour;

RECOMMENDATION

Liaise with local media to encourage reference to and use of guidelines for the reporting of suicide through a summit with local press and media organisations, and to provide information to professionals on the sensitive reporting of suicide. By 20 July 2017.

6. Support research, data collection and monitoring.

RECOMMENDATION

Develop working relationships between individual coroners and local public health teams ensure local plans are evidence based and responsive. Coroners can be invited to become formal members of any local suicide prevention groups or networks .

Coroners can also inform the local authority or Director of Public Health if they identify particular areas of concern, e.g. locations used for suicide, possible clusters of suicide, increase in a particular method or new and emerging method of suicide.

Refresh the Berkshire-wide Suicide Audit for deaths in the period 2014-16 to include new categories based on best practice.

Background

Suicide is a devastating event. It is an individual tragedy, a life-altering crisis for those bereaved, and a traumatic event for communities and services. The impacts are immediately and profoundly distressing. We thus need to be sure that in the Clinical Commissioning Groups (CCGs) and Local Authorities in Berkshire, an alliance of stakeholders takes preventive and ongoing action covering the main risks. The 2012 national strategy ('Preventing Suicide in England') sets us two major objectives: reducing the suicide rate in England, and giving better support to people bereaved or affected by suicide. Those objectives are thus given priority in this strategy.

Suicide is not inevitable. Preventing suicides is a complex and challenging issue, but there are effective solutions for many of the individual factors which contribute towards the risk of suicide. Suicide Prevention work is cost-effective when conducted in accordance with evidence of effectiveness, and by working in partnership. Local Government, statutory services, the third sector, local communities and families each have a role to play.

Whilst suicide causes a vast negative wellbeing impact on family, friends, colleagues, and wider contacts, they also have a huge economic impact. The average cost of a single completed suicide of a working age individual in England was estimated in 2012 to be more than £1.5 million. This includes intangible costs (loss of life to the individual and the pain and suffering of relatives), as well as waged and unwaged lost output, public service time and funeral costs. Non-fatal self-harm also has major - potentially avoidable - cost implications for public services, particularly A&E and acute inpatient services and psychiatric follow-up.

Suicides are not inevitable and are a major issue for society as well as being a leading cause of years of life lost. There are many ways in which services, communities, individuals and society as a whole can help to prevent suicides. Government and statutory services have a role to play in building individual and community resilience. Vulnerable people in the care of health and care services can be supported and kept safe from preventable harm. Interventions can be provided quickly when someone is in distress or in crisis and for vulnerable people in the wider community, practical measure such as debt advice services can make all the difference.

Public Health England (PHE) has recently published a guide to local suicide prevention planning (2016). In it, they identify ten things that everyone needs to know about suicide prevention. These are re-produced here in full, and with kind permission of PHE. Simple to follow and understand, these form the basis of raising awareness of suicide prevention across Berkshire services and populations.

10 Things Everyone Needs To Know About Suicide Prevention

1

Suicides take a high toll

There were 4,882 deaths from suicide registered in England in 2014 and for every person who dies at least 10 people are directly affected.

2

There are specific groups of people at higher risk of suicide

Three in four deaths by suicide are by men. The highest suicide rate in England is among men aged 45-49. People in the lowest socio-economic group and living in the most deprived geographical areas are 10 times more at risk of suicide than those in the highest socio-economic group living in the most affluent areas.

3

There are specific factors that increase the risk of suicide

The strongest identified predictor of suicide is previous episodes of self-harm. Mental ill-health and substance misuse also contribute to many suicides. Suicide prevention strategies must consider and link to programmes of early identification and effective management of self-harm, mental ill-health and substance misuse.

4

Preventing suicide is achievable

The delivery of a comprehensive strategy is effective in reducing deaths by suicide through combining a range of integrated interventions that build community resilience and target groups of people at heightened risk of suicide. Directors of public health and health and wellbeing boards have a central role. Their involvement is crucial in coordinating local suicide prevention efforts and making sure every area has a strategy in place.

5

Suicide is everybody's business

A whole system approach is required, with local government, primary care, health and criminal justice services, voluntary organisations and local people affected by suicide having a role to play. Suicide prevention can also be part of work addressing the wider determinants of health and wellbeing.

6

Restricting access to the means for suicide works

This is one of the most evidenced aspects of suicide prevention and can include physical restrictions, as well as improving opportunities for intervention.

7

Supporting people bereaved by suicide is an important component of suicide prevention strategies

Compared with people bereaved through other causes, individuals bereaved by suicide have an increased risk of suicide and suicidal ideation, depression, psychiatric admission as well as poor social functioning.

8

Responsible media reporting is critical

Research shows that inappropriate reporting of suicide may lead to imitative or 'copycat' behaviour.

9

The cost of suicide justifies investment in suicide prevention work

The economic cost of each death by suicide of someone of working age is estimated to be £1.67 million. This covers the direct costs of care, indirect costs relating to loss of productivity and earnings, and the intangible costs associated with pain, grief and suffering.

10

Local suicide prevention strategies must be informed by evidence

Local government should consider the national evidence alongside local data and information to ensure local needs are addressed.

Strategy Aims

In 2014, the seven Berkshire CCGs and six local authority public health teams across Berkshire began work to refresh the suicide audits previously undertaken and to recommend from this a strategy for reducing suicide risk across Berkshire. This strategy is the result of a study of national research and recommendations plus recommendations of many local stakeholders from a range of organisations.

This strategy proposes co-ordinated prevention across all the elements influencing suicide, from the wider determinants of distress and escalating desperation, and poor mental health, through coordinated local preventive action spanning local authority and voluntary services, and primary and secondary care.

The overall aim of this strategy is:

- To outline how partners across the county will work to prevent suicide in Berkshire.
- To outline the governance structure for Suicide Prevention work in Berkshire.
- To make clear how the public, partners and other stakeholders can deliver the actions outlined herein.

The objectives and six priority areas to meet this aim are also drawn from the National Suicide Prevention Strategy - "Preventing Suicide in England" (DH, 2012), and are intended to be met through coordinated multi-agency actions, under the governance of the Berkshire Suicide Prevention Steering Group.

The objectives of this strategy developed from the national strategy are:

- To reduce suicides in Berkshire by 25% by 2020;
- To ensure better support is provided for those bereaved or affected by suicide.

The priority areas of this strategy drawn from the national strategy are:

1. Reduce the risk of suicide in key high-risk groups;
2. Tailor approaches to improve mental health in specific groups;
3. Reduce access to the means of suicide;
4. Provide better information and support to those bereaved or affected by suicide;
5. Support the media in delivering sensitive approaches to suicide and suicidal behaviour;
6. Support research, data collection and monitoring.

These six priority areas have become the golden thread which runs through this strategy and the action plans which support it. These action plans are for the year 2017/18, whilst the strategy is for the years 2017-2020, taking this to the year when the overarching aim to reduce suicide by 10%, as stated in the Five Year Forward View on Mental Health and incorporated into the Sustainability and Transformation Plans produced by groups of CCGs. There are other recommendations around process and which address the overarching aims and/or a combination of the priority areas.


National Context

Nationally available data on suicides can help place local information on suicides in context. From the national references which include rates, trends, risk factors, suicide methods and evidence of what works to prevent suicides, a local approach for suicide prevention can be developed. This section presents the national data on suicides and is intended to be used as a guide to draw comparisons with local data and information from the Berkshire Audit.

The Office for National Statistics (ONS) provides figures on deaths by suicide, available publicly on its website at: www.ons.gov.uk. Data can be downloaded which shows numbers and rates of death by suicide per 100,000 population. Rates are important as they account for the age and size of populations, so it is more reliable when comparing suicide across age groups and areas.

Suicide is defined in England and Wales as a death with an underlying cause of intentional self-harm and/or an injury/poisoning of undetermined intent (ICD10 codes X60-X84 - all ages and Y10-Y34 - for ages over 15 years). There is an assumption that most injuries or poisonings of undetermined intent are self-inflicted and where there is insufficient evidence to prove that the person intended to kill themselves. This assumption however is not applied to children due to the possibility that these deaths were caused by other situations - such as abuse or neglect. For this reason, data on suicides in England only include persons aged 15 years and over for deaths from injury of undetermined intent and may therefore lead to an under-reporting of deaths as a result of suicide in children.

Figure 1: Age-standardised suicide rates by sex, deaths registered between 1981 and 2014, United Kingdom



Source: Office for National Statistics, National Records of Scotland, Northern Ireland Statistics and Research Agency. 2016.


Figure 1 above shows the age standardised suicide rates for the UK since 1981. A generally downward trend in suicide rates was observed between 1981 and 2007, with a decrease from 15.6 to 10.6 deaths per 100,000 population (see figure 1). There has been a slight overall increase in suicide rates since 2007, to 10.8 per 100,000, which is part of an upward trend since 2007 for both sexes.

Suicide continues to be more than three times as common in males. The male suicide rate in 2013 was the highest since 2001. The lowest male rate since the beginning of the data series, at 16.6 per 100,000, was in 2007.

The highest suicide rate in the UK in 2015 was among men aged 45 to 59, at 22.3 deaths per 100,000, slightly lower than the record high seen in 2013. This age group also had the highest rate among women, at 7.3 deaths per 100,000 population.

Between 2000 and 2011, the rate in this age group was the second highest, behind men aged 30 to 44. Since 2007, the rate in the 45 to 59 age group has been increasing.


Figure 2: Age-specific suicide rate, males, deaths registered between 1981 and 2014, United Kingdom



Source: Office for National Statistics, National Records of Scotland, Northern Ireland Statistics and Research Agency. 2016.

Female rates have stayed relatively constant since 2007. In 2015, the age group with the highest suicide rate for females was 45- to 59-year-olds, with a rate of 7.6 deaths per 100,000 population (see Figure 3). This has been the case since 2003. Analysing this data by 5 year age group shows that females aged 50 to 54 have the highest suicide rate at 8.0 per 100,000 population. Between 1981 and 1994, female suicide rates decreased across all broad age groups apart from 10 to 29 year-olds. Suicide rates for women under 60 have remained relatively constant since 2008, and for women aged 60 and over continue to show a broadly decreasing trend, showing the biggest reduction since 1981.

Figure 3: Age-specific suicide rate, females, deaths registered between 1981 and 2014, United Kingdom



Source: Office for National Statistics, National Records of Scotland, Northern Ireland Statistics and Research Agency. 2016.

A time trend analysis in England suggested that the recent recession in the UK could be an influencing factor in the increase in suicides. The study found that local areas with greater rises in unemployment had also experienced higher rises in male suicides (Barr et al 2012). A review by Samaritans (2012) emphasised that middle-aged men in lower socioeconomic groups are at particularly high risk of suicide. They pointed to evidence that suicidal behaviour results from the interaction of complex factors such as unemployment and economic hardship, lack of close social and family relationships, the influence of a historical culture of masculinity, personal crises such as divorce, as well as a general ‘dip’ in subjective wellbeing among people in their midyears, compared to both younger and older people (Office for National Statistics, 2014).

Suicide by mental health in-patients continues to fall, most clearly in England where the decrease has been around 60% during 2004-14. This fall began with the removal of ligature points to prevent deaths by hanging, but has been seen in suicides on and off the ward and by all methods. Despite this success, there were 76 suicides by in-patients in the UK in 2014, including 62 in England. The 2016 National Confidential Inquiry into Suicide and Homicide by People with Mental Illness found that: many people who died by suicide had a history of drug or alcohol misuse, but few were in contact with specialist substance misuse services; more patients who died by suicide were reported as having economic problems, including homelessness, unemployment and debt; and there has been a rise in the number of suicides by recent UK residents, i.e. those who had been in the UK for less than 5 years, including those who were seeking permission to stay. There are twice as many suicides under crisis resolution / home treatment compared to in-patients.

Hanging, strangulation and suffocation account for the largest number of suicides in males, at 60% of the total. In females hanging and drug related poisoning are the joint most frequent methods, at 38%.

Strategic Context

Local suicide prevention planning is the responsibility of local authority public health teams to deliver with clinical commissioning groups (CCGs), health and wellbeing boards and a wider network of partners. Very recent guidance to inform this strategy has been developed by Public Health England (2016) in partnership with the National Suicide Prevention Alliance.

The need to develop suicide prevention strategies and action plans at a local level and which engage with a wide network of stakeholders in reducing suicide is set out in the government's national strategy for England, Preventing Suicide in England, a cross government strategy to save lives (HM Government, 2012). It is also reinforced by the Mental Health Taskforce's report to NHS England, The Five Year Forward View for Mental Health (NHS England, 2016).

Responsibility for suicide prevention action plans and strategy lies with local government through health and wellbeing boards. Local authorities report on the quality and success of initiatives to improve the health and wellbeing of their populations, using national indicators set out in the Public Health Outcomes Framework. Indicators relevant to suicide prevention include suicide rate, self-harm and excess mortality in adults aged under 75 years with serious mental illness

Around half of people who die by suicide have a history of self-harm, and self-harm is a sign of serious emotional distress in its own right. Mental health promotion, prevention and early intervention are essential to help reduce self-harm in the community that does not present to health services. The effective assessment and management of self-harm by NHS services where people do present with self-harm, particularly in Emergency Departments, represents a huge opportunity to reduce repetition of self-harm and future suicide risk. In June 2013, NICE published a new quality standard to improve the quality of care and support for people who self-harm. The Spending Review (2013) committed to every Emergency Department having constant access to mental health professionals and Public Health Outcomes Framework published in (2013) includes the definition of the new indicator on self-harm. This makes clear the priority given to the prevention and management of self-harm across local authority and NHS services.

PROOF

Evidence Base in Suicide Prevention

The Government published its review of the suicide strategy, *“Preventing suicide in England: Two years on Second annual report on the cross-government outcomes strategy to save lives”* (Department of Health, 2015). This section summaries this, the latest evidence and best practice as identified within the report.

Men and Economic Crisis

A recent study found that men in different age brackets had different suicide risks during the recent recession. Those aged between 35-44 years old experienced increased suicide rates corresponding to economic decline. The study also found the halt in the downward trend in suicide rates amongst men aged 16-34 may have begun before the 2008 economic recession (Coope, et al, 2014).

Self-Harm and Alcohol

There was a higher rate of alcohol-related deaths for those presenting at emergency departments with self-harm for both males and females. Local areas need to ensure that those presenting to hospital with self-harm should be assessed for alcohol problems to identify issues early and get treatment (Bergen, et al, 2014). This is in line with NICE guidelines. In the year following self-harm the risk of suicide is raised 49-fold in the year, this increases with age at initial self-harm (Hawton, et al).

Crisis Resolution

Crisis resolution home treatment services have a key role to play in suicide prevention. Approximately 180 suicides each are patients who are under crisis resolution home treatment services, with approximately 80 among in-patients (Hunt, et al; NCISH 2014).

Primary Care Patients

Both frequent attendance and non-attendance at GP surgeries is linked to increased risk of suicide. For young men, non-attendance is a particular risk factor (NCISH 2002-20012).

Discharge Processes

The first 3 months following discharge from a mental health inpatient episode remains a high risk, with the highest risk at 2 weeks discharge. Community Care reforms which recommend a 7 day follow up have shown positive results although progress has stalled recently (Psychiatry Online).

Self-harm in Prisons

There is an association between self-harm and suicide within the prison setting. Prevention and treatment of self-harm should be part of the suicide prevention efforts within prisons (Hawton, et al, 2014).

National Best Practice in Suicide Prevention

These case studies were reported in, “*Preventing suicide in England: Two years on Second annual report on the cross-government outcomes strategy to save lives*” (Department of Health, 2015).

U Can Cope

The U Can Cope film and online resources were designed for people in distress and those trying to support them, to instil hope, promote appropriate self-help and inform people regarding useful strategies and how they can access help and support. They have been endorsed by the International Association of Suicide Prevention: www.connectingwithpeople.org/ucancope

Social Media

Emerging findings from the research study on Understanding the role of social media in the aftermath of youth suicides (COSMOS), commissioned in support of the suicide prevention strategy, are that:

- Suicidal tweeters show a high degree of reciprocal connectivity (i.e. they follow each other), when compared with other studies of the connectivity of Twitter users, suggesting a community of interest.
- A retweet graph shows that users who post suicidal statements are connected to users who are not, suggesting a potential for information cascade and possibly contagion of suicidal statements.
- The reaction on Twitter to the Hayley Cropper Coronation Street suicide storyline was mostly information/support and debate about the morality of assisted dying, rather than statements of suicidal feelings.
- Tweets about actual youth suicide cases are far more numerous than newspaper reports and far more numerous than tweets about young people dying in road traffic accidents. This suggests that suicide is especially newsworthy in social media. In newspapers there is no significant difference between the two types of death, in terms of number of reports per case.

Nottinghamshire Healthcare NHS Trust and Connecting with People

Nottinghamshire Healthcare NHS Trust is implementing an innovative approach to suicide prevention to improve both patient care and clinical governance. The Trust has developed a team of in-house trainers to deliver suicide and self-harm awareness and response training across the Trust. They are also piloting a web-based App to help to consistently record individual assessments. The App is integrated securely within the NHS system and is based on peer-reviewed clinical tools.

Other Trusts are also involved in the pilot of the App in partnership with the social enterprise *Connecting with People*. The approach being taken:

- Is evidence based and uses peer reviewed clinical tools. It combines compassion with sound clinical governance.
- Is proactive, emphasising safe triage and the co-creation of immediate safety plans for all patients with suicidal thoughts, irrespective of risk. It documents evidence on level of risk and actions agreed to mitigate the risks.
- Has been developed by healthcare practitioners, third sector organisations and service users, for delivery to health and social care practitioners in primary and secondary care and in third sector organisations.

It is an example of how the third sector can work effectively in partnership with the NHS to improve patient care in specialist areas, forming part of the RCPsych OnSite training.

Safety Collaboratives in Mental Health

The South West of England has had a safety collaborative in Mental Health since 2010. This work spread across the South of England in 2013. It involves the majority of Mental Health Trusts in the region. Work streams include getting medicines right, improving physical health care and delivering safe and reliable care. This includes reducing absence without leave from inpatient units and reducing suicides.

North Essex Partnership University NHS Foundation Trust and Samaritans

Three Samaritans branches together with the North Essex Partnership University Foundation Trust have signed a partnership agreement to develop a range of opportunities for patients and staff to benefit from the knowledge, experience and complementary services offered by Samaritans in support of emotional wellbeing and suicide prevention. This will include:

- NHS staff organising for a patient (with their consent) to receive a call from Samaritans.
- Training for non-clinical NHS staff on handling challenging contacts, suicide awareness and emotional well-being.
- Establishing referral pathways between Samaritans and GPs in the area.
- Samaritans presence in Emergency Departments.

This partnership is an example of how the voluntary sector and NHS can deliver better support to people by working together more closely.

Healthtalk Online for parents whose child is self-harming

Drawing on research with families, this website enables parents to see and hear parents and other family members of young people who self-harm share their personal stories on film. The films cover issues such as why young people self-harm, discovering that a young person is self-harming, how they helped their young person, living with self-harm, support and treatment, and what helped them cope.

www.healthtalk.org/peoples-experiences/mental-health/self-harm-parents-experiences/topics.

Staying safe if you're not sure life's worth living

A new online resource developed by the Royal College of Psychiatrists, Connecting with People, Samaritans, Grassroots Suicide Prevention, State of Mind, leading academics, people with lived experience and their carers.

Staying safe if you're not sure life's worth living includes practical, compassionate advice and many useful links for people in distress: www.connectingwithpeople.org/StayingSafe.

Local Context

In Berkshire, the trends in suicides broadly reflect the national trends, and the results from the most recent local suicide audit, carried out in 2015, are shown below.

Of note:

- More males completed suicide than females
- 70% of the deaths recorded between 2007-2014 were in age group 30-59 years
- The percentage of deaths among the unemployed rose from 13% in 2007 to 38% in 2014
- The most common method for suicide was hanging/strangulation.

Local Suicide Audit Results

During 2015, Public Health Teams in Berkshire undertook an audit of suicide and undetermined deaths during the 2012-2014 period. This audit provides an analysis of the most recent audit and includes comparative data from previous audits. The audit defined suicide as a death where the coroner has given a verdict of suicide (based on evidence that the intent was to cause death or take own life) or where an open verdict was reached in a death from injury or poisoning. The definition comprises suicides and open verdicts coded as ICD10 X60-X84 and Y10-YY34.

Data for the audit was collected from Berkshire Coroner Office case notes for people who died from suicide or undetermined injuries during 2012-2014. The audit only includes Berkshire residents who died in the County. It is important to note that there can be a substantial delay between the date of death and the date of registration for suspected suicides, so it is likely that not all deaths from 2014 were included in this audit.


The analysis of suicide data is based on small numbers and is likely to show differences over time or between different groups that are due to random occurrence. However, the analysis of this data can give some indication as to local patterns in suicide deaths. Data from the audit is presented as averages over a three year period to reduce some of the random variations that occur when analysing small numbers. For confidentiality reasons, figures under 5% have been suppressed and data is shown at a Berkshire level, rather than by individual local authorities.

120 deaths were included in the Berkshire suicide audit for 2012-14. 70% of these were classified as suicide by the coroner and the other 30% were undetermined deaths / open verdicts.

Gender

Data from all recent audits show that males have a higher suicide rate compared to women in Berkshire. This is consistent with national figures.

Figure 4: Suicide and undetermined deaths in Berkshire by gender (2007-09 to 2012-14)



Age

70% of the deaths recorded in 2012-14 were for people aged 30-59. The audit was carried out using the age brackets as below, and not the age brackets used by the Office for National Statistics (ONS). Future audits will use the ONS brackets to achieve comparison between local data and national data.

Age group	2012-2014
10-19	*
20-29	13%
30-39	23%
40-49	23%
50-59	24%
60-69	*
70-79	*
80-89	7%

Ethnicity

The majority of people dying from suicide or an undetermined death in Berkshire are White-British. This is largely representative of Berkshire's population. In 2011, 73% of Berkshire's population were from a White-British background, ranging from 35% in Slough to 80% in West Berkshire. The majority of deaths from other ethnic groups (Asian/Asian-British and White-Other) were Slough residents and this also reflects the Borough's population profile. It is important to note that 15% of the cases included in the 2015 audit did not have an ethnic origin recorded in the audit. This is a higher proportion than the previous audit and will therefore have affected the validity of the analysis for 2012-2014 data.

Ethnicity	2007-2009	2008-2010	2009-2011	2012-2014
White-British	77%	75%	77%	61%
White-Other	10%	15%	13%	13%
Asian/Asian-British	<5%	<5%	<5%	12%
Black/Black-British	<5%	<5%	<5%	0%
Not Known	<5%	<5%	<5%	15%

Diurnal and Seasonal Variation

The following tables illustrate the day of the week and seasons in which deaths from suicide occurred in Berkshire during 2007-2011 and 2012-2014.

Day of the week	2007-2009	2008-2010	2009-2011	2012-2014
Monday	19%	21%	21%	20%
Tuesday	16%	17%	17%	13%
Wednesday	16%	11%	10%	9%
Thursday	10%	10%	15%	16%
Friday	8%	9%	7%	14%
Saturday	19%	14%	15%	13%
Sunday	14%	17%	15%	13%

The data shows a relatively even spread across the whole week, with no particularly 'common' day.

Season	2007-2009	2008-2010	2009-2011	2012-2014
Winter (Dec-Feb)	24%	23%	27%	28%
Spring (Mar-May)	29%	30%	27%	31%
Summer (Jun-Aug)	25%	21%	21%	21%
Autumn (Sept-Nov)	21%	26%	28%	18%

Figure 5: Suicide and undetermined deaths in Berkshire by day of week (2007-09 to 2012-14)






Figure 6: Suicide and undetermined deaths in Berkshire by season (2007-09 to 2012-14)



Marital and Living Status

Recent data from the Office for National Statistics shows that 13% of usual residents in England and Wales were living on their own in 2011. The table below indicates that those living alone in Berkshire are therefore over-represented in suicide deaths. This percentage has reduced from 49% in 2009-2011 to 34% in 2012-2014, however it is still the main living status recorded. It is important to note that the number of people with a living status not recorded or not known is higher in 2012-2014 (13%), which makes comparisons of data difficult.

Figure 7: Suicide and undetermined deaths in Berkshire by living status (2007-09 to 2012-14)



The table below shows that there were more deaths from single people in the two audit periods, ranging from 39% to 45%.

Marital status	2007-2009	2008-2010	2009-2011	2012-2014
Single	45%	39%	39%	40%
Married	23%	29%	30%	29%
Divorced	14%	13%	13%	8%
Separated	10%	7%	7%	<5%
Widowed	4%	6%	7%	<5%
Co-habiting	<5%	<5%	5%	10%
Not stated	<5%	<5%	<5%	6%

Employment Status

Some studies have indicated that there is a strong independent association between suicide and individuals who are unemployed (Lewis and Sloggert, 1998). Unemployment in the Thames Valley is low, although there has been some fluctuation between 2007 and 2014. The lowest level of unemployment during this time was 3.4% in Jul-07 to Jun-08, with the highest rate of 6.1% in Apr-09 to Mar-10.

Data from the 2012-2014 Berkshire audit shows that 38% of people dying from suicide and undetermined deaths were unemployed. This is an over-representation of the population, considering that only 4-5% of people were unemployed during that time period. This is also a notable increase on the figures from 2007-2011, which ranged from 11%-14%. This change may be down to a random occurrence, due to small numbers.

Employment status	2007-2009	2008-2010	2009-2011	2012-2014
Full Time	46%	51%	55%	36%
Part Time	5%	<5%	<5%	<5%
Unemployed	13%	11%	14%	38%
Student	6%	6%	<5%	<5%
Retired	18%	17%	17%	11%
Long-term illness/ disability benefits	<5%	<5%	<5%	<5%
Housewife/husband	<5%	<5%	<5%	<5%
Not known	8%	5%	<5%	12%

Suicide Note

The table below shows the proportion of deaths where a suicide note was left.

Left a suicide note?	2007-2009	2008-2010	2009-2011	2012-2014
Yes	29%	32%	40%	36%
No	71%	68%	60%	54%
Not known	0%	0%	0%	10%

Housing Status

A large number of the cases included in the 2012-2014 audit did not capture the housing status for people, which means that the data cannot be presented in this analysis.


Housing status	2007-2009	2008-2010	2009-2011	2012-2014
Owner/Occupier	46%	46%	52%	35% of these cases did not have a housing status recorded and therefore this data cannot be presented
Privately Renting	41%	33%	25%	
Council House/ Housing Association	5%	9%	11%	
With Parents	<5%	<5%	<5%	
Supervised Hostel	<5%	<5%	<5%	
Unsupervised Hostel	<5%	<5%	<5%	
Other	<5%	<5%	<5%	
Not Known	<5%	<5%	<5%	

Location of event

The majority of deaths identified in the local audits took place in the person's own home or another person's home. This proportion has continued to increase from 62% in 2007-2009 to 71% in 2012-2014.

Location of event	2007-2009	2008-2010	2009-2011	2012-2014
Own/Others Home	62%	64%	66%	71%
Railway Station/ Crossing	6%	9%	9%	<5%
Stream/river	<5%	9%	8%	6%
Other	28%	18%	17%	13%
Not known	0%	0%	0%	6%

Figure 8: Suicide and undetermined deaths in Berkshire by location of event (2007-09 to 2012-14)



Methods Used

Suicide methods can be classified as either 'active' or 'passive'. Active methods are quick and effective allowing little time for reconsideration. Such methods are hanging, shooting, jumping in front of a train or from a height. Among the general population hanging, strangulation and suffocation has been identified as the most common cause of suicide for men. Passive methods are less violent and allow some time for reconsideration or intervention (e.g. self-poisoning, carbon monoxide). Hanging/strangulation has been the most common cause of death over 2007-2014.

Methods used	2007-2009	2008-2010	2009-2011	2012-2014
Hanging / Strangulation	54%	47%	48%	49%
Carbon Monoxide Poisoning	8%	<5%	<5%	<5%
Jumping / laying before a train	6%	9%	9%	<5%
Jumping from a height	11%	11%	8%	<5%
Self-Poisoning	10%	9%	12%	0%
Drowning	<5%	7%	7%	6%
Other	7%	12%	14%	38%
Not known	0%	0%	0%	<5%

Alcohol and drugs taken at time of death

The audit of people dying from suicide and undetermined deaths during 2012-14 identified whether alcohol or prescribed drugs were detectable in the deceased. This data was not collected in the previous audit. The tables below show that at least 36% of people who died in 2012-14 had taken alcohol prior to their death and at least 42% had taken prescribed drugs, and outlines those drugs that were implicated in suicide deaths.

Alcohol present?	2012-2014
At intoxicating level	23%
At non-intoxicating level	13%
No alcohol detected	54%
Not known	11%
Prescribed drugs present?	2012-2014
At fatal level	14%
At intoxicating level	8%
At therapeutic level	20%
No prescribed drugs detected	43%
Not known	16%

Drugs implicated	Male	Female
Antidepressants	✓	✓
Paracetamol	✓	
Coproxamol or similar	✓	✓
Benzodiazepine	✓	
Other hypnotic		
Anti-psychotic	✓	✓

Other substances implicated in suicide deaths in 2012-14 were:

Other substances	Male	Female
Amphetamines	✓	✓
Ecstasy	✓	
Crack/Cocaine	✓	
Ketamine	✓	
Heroin	✓	✓
Opiates	✓	
Methadone	✓	✓

Personal, Social and Health Factors associated with deaths from suicide

The following factors were identified from records at the Coroner's Office as being associated with suicide:

Factor identified	2007-2009	2008-2010	2009-2011	2012-2014
Relationship problems	14%	6%	<5%	29%
Financial problems	9%	6%	<5%	24%
Depression	25%	42%	51%	67%
Low self esteem	<5%	<5%	<5%	Not collected
Other Mental health Issues	8%	8%	<5%	Not collected
Pending Police Investigation	<5%	<5%	<5%	12%
Family bereavement	<5%	<5%	<5%	12%
Physical Health	8%	<5%	<5%	33%
Job related	<5%	<5%	<5%	17%
Not Stated	15%	13%	20%	-

2015 Data Update

The most recent raw data on the number of suicides in Berkshire was released in December 2016 by the Office for National Statistics for the year 2015. This shows an increase across Berkshire as a whole rather than a small decrease as seen in England and the South East. Caution should be employed as these raw data do not give the detail required to indicate trends or draw conclusions.

	2014	2015	Difference
Bracknell Forest	5	10	+ 5 (+ 100%)
RBWM	11	11	0
Slough	15	9	- 6 (- 40%)
Reading	12	18	+ 6 (+ 50%)
West Berkshire	5	6	+ 1 (+ 20%)
Wokingham Action	6	14	+ 8 (+ 233%)
Berkshire Total	54	68	+ 14 (+126%)
SE England Total	794	756	- 5%
England Total	4882	4820	- 1%

Local Governance Structures

In order to facilitate the production of this strategy and to steer the Berkshire-wide audit of suicides, a strategic group was convened with representatives from organisations across county. This worked under the identity of the Berkshire Suicide Risk & Self Harm Reduction / Prevention Steering Group. During 2015/16 as key staff changed, the group has lost some of its membership and had become less strategic. The original terms of reference state that the group:

“will provide public health leadership and advice to support a joint approach to achieve real change in the prevention of suicides and self-harm through actions taken by member organisations. It will facilitate the bringing together of clinicians, professionals and organisations, with the patient’s voice, to deliver surveillance data to support projects / programmes to prevent suicides and offer support to those who are bereaved.”

Public Health England (2016) suggests that the membership of suicide prevention partnerships is made up of representative working with adults, children and young people. The following diagram suggests the range of partners who may be included.



The Berkshire Steering Group will need to own this strategy, and the membership should be updated to ensure a closer fit with the groups suggested above. The membership of the current group as at December 2016 is detailed in Appendix 8.

RECOMMENDATION

That the Berkshire Steering Group re-visits their terms of reference and membership and become known as the “Berkshire Suicide Prevention Steering Group”, with the aim of providing the governance to this strategy and its action plans.

There are other groups for whom this strategy and its action plans are an important issue, and these include commissioning boards in the East Berkshire and West Berkshire Confederations of CCGs; Health and Wellbeing Boards; Health Overview and Scrutiny Committees; Adult Safeguarding Boards; and Community Safety Partnerships. In order to get full endorsement of this strategy and for organisations to commit to their action plans, the terms of reference should ensure that the links to these other structures are robust and transparent.

Members of the Steering Group could be asked to act as suicide prevention champions. These are individuals who get involved in specific pieces of suicide prevention work - and might include people who have been bereaved by suicide or those with a special interest or expertise. They can be pivotal in raising issues regarding suicide awareness locally, and drive forward the action plans of their agencies. A specific initiative to engage the elected members of councils as Mental Health Champions may provide an opportunity for them to also speak out on suicide prevention. Details of this initiative are available here: <http://www.mentalhealthchallenge.org.uk/the-challenge/>

RECOMMENDATION

That organisations represented on the Berkshire Suicide Prevention Steering Group consider nominating a suicide prevention champion from within their membership.

Monitoring & Evaluation and Progress

This is a pan-Berkshire and multi-agency Strategy and the action plans for this strategy which are overseen by the Berkshire Suicide Prevention Steering Group are set out below.

Individual Borough action plans for each of the six Berkshire Unitary Authorities are also included and are set out in appendices 2-7. These give a more local set of priorities and respond to the particular geographical issues, population structures and general health needs of the Authorities.

Other agencies which are part of the Steering Group may have their own action plans and an objective of this strategy is to bring these into one combined action plan as far as possible and to share openly the actions plans of all agencies in order to learn from one another; to avoid unnecessary duplication of effort or resources; and to encourage co-production of outcomes.

Links to Other Local Strategies

This is the first comprehensive Berkshire-Wide Suicide Prevention Strategy and action plans have been produced for the year 2017-18. One of the objectives is to ensure that this strategy, its aims and objectives are shared and upheld in the strategies, action plans and objectives of all those groups across Berkshire who are committed to improving health outcomes, promoting wellbeing, removing the stigma associated with mental health and preventing suicides.

Local Joint Health and Wellbeing Strategies and their action plans should endorse this Strategy and Health and Wellbeing Boards are key to the governance of this Strategy and the Steering Group. Through tightly-knit joined up thinking, organisations, individual and communities across Berkshire can come together to make the progress necessary to reduce suicides in our populations.

RECOMMENDATION

That all agencies represented on the Steering Group commit to an annual Action Plan of their own which can then be brought together to create a Berkshire-wide Action Plan for each year of the Strategy.

Local Best Practice in Suicide Prevention

Thames Valley Suicide Prevention and Intervention Network (SPIN) CALMzone

The Campaign Against Living Miserably (CALM) was originally a Department of Health helpline project on suicide prevention particularly targeting younger men using marketing methodology and images to specifically engage with this audience on issues surrounding mental distress and social alienation. The resources produced directed men to a special helpline, and latterly to web-based resources. In 2000, a partnership of six areas in the North-west of England commissioned this work for young men in Merseyside, which continued when CALM transferred into a national charity. There is a local CALMzone Coordinator who promotes CALM across Merseyside in collaboration with the local community - pubs and clubs, venues and universities, sports teams and clubs - to encourage them to join and promote the campaign.

In 2015, the local authorities across the Thames Valley through SPIN funded a Thames Valley CALMzone, and employed a coordinator undertaking similar promotions as in Merseyside. CALM have provided local commissioners with anonymised reports on numbers and trends of calls and web chats across the Thames Valley. As well as providing funding to support the helpline, the commissioners ensure CALM has an up-to-date local database of agencies which local callers can be referred to. Berkshire local authorities have continued to fund the helpline until June 2017, although the local coordinator post is no longer funded.

RECOMMENDATION

Evaluate the Berkshire CALMzone and recommission targeted suicide prevention work for younger men and middle aged men.

Real Time Suicide (and near fatal self-harm) Surveillance

It is important to have a real time overview of self-inflicted deaths/suspected suicides and near fatal self-harm in order to provide timely support for those bereaved and affected, pick up community risks of contagion or suicide clusters and identify public places where suicides/incidents of near fatal self-harm appear to occur with increasing frequency. All of these activities contribute to suicide reduction and prevention in line with national and local strategy. Thames Valley Police (TVP) and the Thames Valley Suicide Prevention and Intervention Network (SPIN), supported by funding from the Thames Valley Strategic Clinical Network are collaborating to build on the supportive signposting for people bereaved by suicide work and develop a robust real time surveillance process.

In simple terms this process is as follows:

- TVP identifies and collates suspected suicides on the Gen 19 sudden death form.
- Coroner's officers send Gen 19s of suspected suicides to a central TVP email for monitoring.
- Details of the incidents in real time are thereby collated and are available for analysis, reporting and provide the ability to respond.
- Details of families who consent to 'Supportive Signposting' are sent to a central NHS England suicide bereavement address.
- Supportive literature and referral signposting links to organisations and charities are provided to relatives.

RECOMMENDATION

Ensure bereavement information and access to support is available to those bereaved by suicide, including professionals involved in the case.

Further data is being collated from the following sources;

- NHS England monitor the strategic executive information system database for suspected suicides and near fatal self-harm.
- Links are being established with British Transport Police to monitor suspected suicides and near fatal attempts on the railways.
- Links are being established with prisons to monitor prison suspected suicides and near fatal attempts.
- Links are to be established with the general hospital psychiatric liaison services to monitor incidents of near fatal self-harm.

All of this information will be reviewed by the TVP and SPIN leads and figures and concerns will be communicated to local public health suicide prevention leads for consideration within the local multi-agency suicide prevention action groups. A hub of SPIN comprising TVP, public health, NHS and the University of Oxford Department of Psychiatry Centre for Suicide Research has been established to maintain oversight of the regional prevalence of suicide with the aim of collaborating where indicated, in order to respond to issues that concern the whole geography, for example contagion and clusters.

Berkshire Healthcare Foundation NHS Trust (BHFT) Zero Suicide Programme

Berkshire Health Care Foundation NHS Trust have been inspired by the pioneering zero suicide approach within the Henry Ford Hospital System (USA). The Henry Ford Hospital System managed to implement a philosophy and practice of 'perfect depression care' which led within four years to a 75% drop in suicides, and eventually to years without a single suicide. For BHFT Zero suicide means using the ambitious target of zero to help focus on this quality improvement issue. Thinking in this way encourages tracking of best practices and formally incorporating them into how service users are treated. Most importantly, it encourages Trust staff to work with in collaboration with service users, carers and primary care colleagues to focus on genuine service user engagement and identification of need.

Based on their analysis of research and Trust data showing key patterns and risk points for suicide, BHFT have set key priority areas for their zero suicide campaign.

1. Optimise systems to enable staff to focus on engagement and collaborative approaches to risk assessment and management with service users and carers at the centre.
2. Training and supervision to equip staff with skills and competence to practice recovery focussed approaches to suicide risk and enable positive risk management and safety planning.
3. Development of a BHFT suicide surveillance dashboard - real time information on service gaps and analysis of patterns to inform practice and training.
4. Collaborating with colleagues, service users and carers as part of a wider suicide awareness campaign beyond secondary mental health care.

Royal Berkshire NHS Foundation Trust (RBFT) Zero Suicide Programme

Royal Berkshire NHS Foundation Trust is working in partnership with BHFT services, Thames Valley Police and Samaritans (Reading) to support the BHFT's ambitious target of zero suicide.

Based on their analysis of research, Trust data showing key patterns and risk points for suicide and patient, family and carer experience RBFT have set priorities for their zero suicide plan which is driven and monitored through a multiagency Suicide and Self Harm Prevention Governance Group chaired by their Mental Health Coordinator related to:

- 1.** Collaborative working with the Psychological Medicine Service (PMS) and patient families and carers to risk assess individuals who attend in crisis.
- 2.** Environment, Estate and capacity of teams.
- 3.** Training, supervision and support to provide staff with skills and competence to recognise risk and manage it proactively in partnership.
- 4.** Collaborative working with multiagency colleagues, patients, families and carers and our staff as part of a wider 'Let's Talk Mental Health' campaign.

DIGITAL
SOLUTIONS
PROOF

Areas of High Frequency

Due to their geography, design or operational use, there are places which present easier access to the means of suicide than others. This could be as a result of their isolation from staff operating their functions; because they are more generally isolated from crowds and the general public; or because life-threatening hazards exist which are generally mitigated by normal operation. They may have become known as places where suicides have occurred previously, either via media reports, or word of mouth.

The Railway Network

The railway network, mostly operated by Network Rail, is in places associated with higher frequencies of suicides, injurious attempts at suicide and suicide attempts and other incidents of people in hazardous positions which do not cause physical injury. The rail network in Berkshire includes a section of the main Great Western Railway routes from London to Wales and the South West, as well as sections of suburban rail lines and minor branch lines. The Great Western lines feature high-speed trains, and is presently being electrified by means of overhead cables. Most of the suburban rail lines are electrified using a third rail system. As well as a high volume of passenger trains, most local lines also feature freight trains operating throughout the day and night.

On average there are 255 suicides on the network per annum. Rail staff particularly drivers, are likely to be severely traumatised by these events and some may never return to work and therefore might need to access support services because of that. Network Rail operates a comprehensive programme of suicide prevention, working to reduce the potential for suicides to occur on the rail network and the industry sees its potential as going beyond that by seeking to do all it can to prevent suicides in its neighbouring communities. In 2015 Network Rail, together with British Transport Police, and Samaritans agreed a process whereby any location that experienced three suspected suicide or injurious attempt incidents (or a combination of the two) would be subject of an escalation process. This would mean that enhanced working would be taken by all three parties in order to prevent further incidents at that that location. In Berkshire, there are locations where this process has been enacted. Actions taken at these locations include engineering solutions, such as the replacement of crossing with overbridges; or the fencing off of platforms on non-stopping fast lines; and the placement of Samaritans posters across the location.

RECOMMENDATION

[That local authority public health teams take the leadership for liaison with any “Escalation Process” in their area, and report on progress to the Steering Group.](#)

The Motorway and Roads Network

Most motorways and trunk roads (the strategic road network) are the operational responsibility of Highways England, with most other roads being the responsibility of local authorities, whilst some roads and byways are in private ownership. In Berkshire, the main London to South Wales Motorway, the M4 passes through the length of the county through all of the six unitary authorities and is managed by Highways England, whilst the adjoining A329(M) motorway is the responsibility of Wokingham Borough Council. The speed of traffic on these roads and other major roads together with their overbridges provide places which can give access to the means of suicide. In 2013, the Highways Agency Traffic Officer Service attended 293 of 652 suicide/attempted suicide incidents on the strategic road network. Between April 2013 and December 2014, there were over 1,500 incidents which were brought to the attention of a Traffic Officer

(Sutherland, 2015). Definitions in this area are not clear, but this does seem to indicate an increase in the reporting of road network incidents, if not in the number of incidents themselves.

RECOMMENDATION

That the Steering Group seek a named Highways England officer to act as a liaison link and group member, and to share real-time intelligence of highways network incidents.

Car Parks and Tall Buildings

Berkshire towns feature a number of multi-storey car parks of which some are in the management or ownership of the Local Authorities. There are also many other tall buildings both residential and commercial, and together these locations can sometimes provide an access route to a means of jumping from a height.

The Horsham branch of Samaritans is working with a local shopping centre and car park where suicides have occurred. They offer sessions for the car park attendants who are generally the first on the scene. They also have an arrangement with the shopping centre to call Samaritans if there is an incident, either to support the staff involved or to support shoppers/shop workers more generally if the incident was widely witnessed. (Sutherland, 2015). This is a simple intervention in which suicide prevention training could be incorporated.

Local Authority Settings

Local authorities may be responsible as owners, operators or managers of other facilities and locations where suicides may take place. This may be because of their isolation or due to their inclusion of specific means of suicide within them. Generally the local authorities in Berkshire look after many hectares of open space; parkland; and woodland, some of which may be managed as part of the highways network; but with most likely to be part of an open spaces portfolio. There is also significant waterside public realm managed or owned by the authorities. The risks at these sites include strong, tall trees as a means of hanging; access to water features such as lakes, rivers and canals which pose a risk of drowning; and dense undergrowth which could allow a person to die through neglect and exposure. Council staff and contractors may have an enhanced role to play in identifying suicide risks and in supporting people who appear to be in distress.

RECOMMENDATION

That local authority public health teams work with other council departments such as car parks and open space services to identify local actions to prevent suicide including staff awareness training.

Mental Health Crisis Care Concordat

The Mental Health Crisis Care Concordat is a national agreement between 22 national agencies involved in the care and support of people in crisis and includes health, policing, social care, housing, local government and the third sector. It sets out how these agencies will work together better to make sure that people get the help they need when they are having a mental health crisis. Local areas have submitted declarations and developed action plans for the improvement of local mental health crisis care for their areas.

The Concordat focuses on four main areas:

- Access to support before crisis point - making sure people with mental health problems can access help 24 hours a day and are taken seriously.
- Urgent and emergency access to crisis care - making sure that a mental health crisis is treated with the same urgency as a physical health emergency.
- Quality of treatment and care when in crisis - people are treated with dignity and respect, in a therapeutic environment.
- Recovery and staying well - preventing future crises by ensuring that people are referred to appropriate services.

Although the Crisis Care Concordat focuses on the responses to acute mental health crises, it also includes a section on prevention and intervention. This strategy includes and reflects some of the key messages and recommendations from the concordat, aiming to reinforce a commitment by partners to work together in preventing and managing crises.

The local Crisis Care Concordat has, common to this strategy, been set at Berkshire-wide level, and has a comprehensive action plan, and certain actions include specific suicide prevention actions. These relate to the work of British Transport Police and the escalation process and staff training issues. They are more detailed than the recommendations and actions set out in this strategy, but there is strategic fit. There is a need to ensure full reference to this strategy in the Crisis Care Concordat action plans, and for further synergies to be explored.

RECOMMENDATION

Ensure strategic fit with the action plans of the Berkshire Crisis Care Concordat.

Gap Analysis and Emergent Berkshire-Wide Concerns

A gap analysis was undertaken by members of the Steering Group to identify the areas of the National Strategy which were not seen to be adequately addressed across Berkshire, taking into account the results of the local Suicide audit and the demography of the six unitary authorities. Some emergent concerns have also been captured which reflect discussion on the audit findings.

High Risk Groups

This strategy recognises that individuals may fall into two or more high-risk groups. Conversely, not all individuals in the group will be vulnerable to suicide. Other risk factors, such as loneliness, social circumstances and physical illness, must also be considered within the wider context or risk (Preventing suicide in England, Department of Health, 2012).

Berkshire is home to the University of Reading and other higher and further education establishments. Although the risk of suicide in the student population locally has not been established, recent ONS data (ONS, 2012) has shown a substantial increase nationally in both male and female suicides in the student population from 2007-2011.

Carers and people with long-term conditions have been highlighted as a local population at particular risk and this has been reinforced by the investigations into domestic homicides where a partner had subsequently taken their own life or attempted to. Adult Social Care and Public Health Outcomes Frameworks record measures of carer social interaction and that of people receiving care which give an insight into the vulnerabilities of these groups, and these are highlighted in the PHE Suicide Prevention Profiles. Not all people with a long-term health condition will be captured within these data, however; and the impact of symptoms such as chronic pain and reduced mobility, and access to certain medicines make this a group with heightened risk and access to means of suicide.

Berkshire no longer contains a prison. People in the criminal justice system will be imprisoned in neighbouring counties, which could make access visits more difficult for family and friends leading to increased isolation for the imprisoned.

Self-Harm continues to be an important risk factor for suicide and growing evidence to support using self-harm as an outcome measure for suicide prevention work with evidence showing that hospital presentation following self-harm is a clear risk factor for suicide (Hawton et al. 2012). There are around 200,000 episodes of self-harm that present to hospital services each year in England, although the true scale of the problem is not known as many people who self-harm do not attend A&E, or seek help from health or other services. Around 50% of people who die by suicide had a history of self-harm, in many cases with an episode shortly before their death, and around 15% of those who die by suicide have carried out an act of self-harm leading to presentation at hospital in the year before their death (PHE 2016).

The table below shows the rates of self-harm and suicide in the six authorities in Berkshire from the PHE Suicide Prevention Profiles (PHE, 2016A). All authorities have lower rates than England, although there is quite some variation across the authorities. It is important to ensure implementation of the NICE standards and pathways CG16 and CG133 for managing patients who self-harm.

Indicator	Period	England	SE England	Bracknell Forest	Reading	Slough	West Berkshire	Windsor & Maidenhead	Wokingham
Hospital stays for Self-harm	2014-15	191.4	193.1	118.3	130.0	162.2	127.0	150.6	91.1
Suicide Rate persons	2013-15	10.1	10.2	8.1	11.0	8.8	7.0	7.1	6.0
Suicide rate (male)	2013-15	15.8	15.9	*	19.0	14.8	*	*	*
Suicide rate (female)	2013-15	4.7	4.8	*	*	*	*	*	*

Source: PHE Prevention Profiles. 2016

RECOMMENDATION

Implement the NICE guidelines on self-harm, specifically ensuring that people who present to Emergency Departments following self-harm receive a psychosocial assessment.

Tailor approaches to improve mental health in specific groups

Work on detailed Mental Health Strategies is underway across the Berkshire East and Berkshire West health systems. It will be important to ensure a good strategic fit between this strategy and those that are developed. Mental health and wellbeing promotion will remain important objectives of both strategies.

RECOMMENDATION

Work to provide and commission interventions which improve the public's mental health. These may include: awareness of mental health and peer support in young people; anti-bullying campaigns in schools; addressing stigma and social isolation in older people; workplace health promotion and support with local business; working with police on mental health literacy; and addressing issues relevant to the local population.

Support research, data collection and monitoring

With real-time surveillance giving information on suicides and many near fatal self-harm events, there is concern that not all events will be recorded, for instance those attempted suicides which occur on the highways network.

There is further analysis of Coroner's case notes that is recommended as good practice, such as the last contact with a GP which have not been captured in the last local audit. A new audit should be run with this new category for deaths in the period 2014-16, beginning as soon as practicable. This can then be appraised alongside data received through real-time surveillance; gaps identified and protocols and policies put in place to ensure that data can be confidentially shared for the purposes of identifying trends and clusters in order to take appropriate preventative actions.

RECOMMENDATION

Develop working relationships between individual coroners and local public health teams ensure local plans are evidence based and responsive. Coroners can be invited to become formal members of any local suicide prevention groups or networks.

Coroners can also inform the local authority or Director of Public Health if they identify particular areas of concern, e.g. locations used for suicide, possible clusters of suicide, increase in a particular method or new and emerging method of suicide.

Refresh the Berkshire-wide Suicide Audit for deaths in the period 2014-16 to include new categories based on best practice.

Domestic Homicide Reviews

Domestic Homicide Reviews (DHRs) are part of the Domestic Violence, Crime and Victims Act 2004 and became law as of the 13th of April 2011. They do not replace but are in addition to an inquest or any other form of inquiry.

DHRs are one way to improve responses to domestic violence and aim to prevent the avoidable death of a member of the community. The review helps to ensure that public bodies including health, local authorities, police and other community based organisations understand the factors surrounding the death and identify where responses to the situation could have been improved. From this, the agencies involved are in a stronger position to learn appropriate lessons, including those involving joint working. A DHR does not seek to lay blame but to consider what happened and what could have been done differently. It also recommends actions to improve responses to domestic violence situations in the future.

DHRs are commissioned by Community Safety Partnerships where a death of a resident has occurred in accordance with the criteria set out in the Home Office Multi Agency guidance;

‘Domestic homicide review means a review of the circumstances in which the death of a person aged 16 or over has, or appears to have, resulted from violence, abuse or neglect by:

- (a) A person to whom he was related or with whom he was or had been in an (a) intimate personal relationship, or
- (b) A member of the same household as himself, held with a view to identifying the lessons to be learnt from the death.’

Updated DHR guidance was published in December 2016 and the DHR process is also now available to cover historic victims of domestic abuse:

“Where a victim took their own life (suicide) and the circumstances give rise to concern, for example it emerges that there was coercive controlling behaviour in the relationship, a review should be undertaken, even if a suspect is not charged with an offence or they are tried and acquitted. Reviews are not about who is culpable.”

Such circumstances are likely to be rare; however the duty to undertake a DHR if required may place additional burden on those implementing suicide prevention locally. However, this must be balanced with the likelihood of new learning, which should be fed back into the Berkshire Suicide audit process.

Berkshire-Wide Action Plan 2017-18

Areas for Action	Specific Risk Groups	Action in 2017-18	Timescale by:	Delivery Lead
Overarching Aims		Sign off / endorsement of this Berkshire Suicide Prevention Strategy by all Health & Wellbeing Boards in Berkshire.	1 April 2017	Lead Consultant Mental Health
		All other local governance completed thus enabling this strategy to be owned by the Local Authorities and their partners.	1 April 2017	Local PH Mental Health Leads
		Launch of strategy at multi-agency suicide prevention summit.	15 Oct. 2017	Strategic DPH
		Identify Suicide Prevention Champions to promote the Berkshire Strategy; and to work within the Local Authorities and partner agencies.	15 Oct. 2017	Local PH Mental Health Leads
		Develop Berkshire-wide information sharing protocols to best utilise real time surveillance of suicides and near misses, in order to respond promptly to local trends and risks to reduce risk of clusters, and inform future service delivery.	30 July 2017	Lead Consultant Mental Health
		The Steering Group revisit their terms of reference and membership and become known as the "Berkshire Suicide Prevention Steering Group", with the aim of providing the governance to this strategy and its action plans.	1 April 2017	Steering Group Members
		Ensure strategic fit with the action plans of the Berkshire Crisis Care Concordat.	1 April 2017	Lead Consultant Mental Health
National Strategy				
1. Reduce the risk of suicide in key high-risk groups	Men	Evaluate the Berkshire-Wide CALMzone initiative and agree Berkshire-wide commissioning of specific support services for men for future years. To include future commissioning of CALMzone for younger men; and services for middle aged men and older men.	15 Oct. 2017	Lead Consultant Mental Health

National Strategy				
	People who self-harm	Ensure agencies have plans to Implement the NICE guidelines on self-harm.	15 Oct. 2017	Lead Consultant Mental Health
	People who misuse substances	Ensure local strategies and contracts for DAAT services include suicide prevention objectives.	Ongoing work	Local PH Mental Health Leads
	People in mental health care	Support BHFT in its Zero Suicide Approach, and support local prevention work across the care system.	Ongoing work	Steering Group Members
	People in contact with the criminal justice system	Identify local actions to prevent suicide in those in contact with the criminal justice system, recognising increased incidence of self-harm in the prison population.	30 July 2017	Local PH Mental Health Leads
	Occupational groups	Ensure local health trusts and providers can demonstrate actions to prevent suicide and promote mental wellbeing amongst their staff.	30 July 2017	Steering Group Members
		Identify particular local action plans for those in agricultural / land-based industries.	30 July 2017	Local PH Mental Health Leads
2. Tailor approaches to improve mental health in specific groups	Community based approaches	For the Steering Group to assess community-based interventions which may be best delivered at scale across the county.	Ongoing work	Steering Group Members
	Suicide prevention training	Coordinate a database on evidence based suicide prevention training programmes and providers across the county.	Ongoing work	Steering Group Members
	People vulnerable due to economic circumstances	For the Steering Group to solicit data from each LA on key indicators that may highlight risk: e.g. number of homelessness presentations.	Ongoing work	Steering Group Members
	Pregnant women and those who have given birth in last year	To undertake a needs assessment of this group in relation to suicide prevention.	30 July 2017	Local PH Mental Health Leads

National Strategy				
	Children and young people	Through LSCBs, identify local actions to prevent suicide in children and young people.	30 July 2017	Local PH Mental Health Leads
3. Reduce access to the means of suicide		<p>Support Network Rail, British Transport police and The Samaritans with local escalation process locations and general suicide prevention work.</p> <p>Investigate suicides on council owned land and properties, and agree a local action plan.</p> <p>Analyse local data gathered from the suicide audit and/or real-time surveillance to identify trends and clusters and take appropriate action(s) e.g. work with local media.</p> <p>The Steering Group seek a named Highways England officer to act as a liaison link and group member, and to share real-time intelligence of highways network incidents.</p>	<p>Ongoing work</p> <p>15 Oct. 2017</p> <p>Ongoing work</p> <p>1 April 2017</p>	<p>Steering Group Members</p> <p>Local PH Mental Health Leads</p> <p>Local PH Mental Health Leads</p> <p>Lead Consultant Mental Health</p>
4. Provide better information and support to those bereaved or affected by suicide		Promote resources for people bereaved and affected by suicide (e.g. Help is at Hand and National Suicide Prevention Alliance resources), and support services such as SOBS (Survivors of Bereavement by Suicide).	Ongoing work	Steering Group Members
5. Support the media in delivering sensitive approaches to suicide and suicidal behaviour		<p>Convene a Berkshire-wide Summit on journalism and reporting standards with local press and media organisations, to develop and agree standards for reporting.</p> <p>Agree a local action plan with the local communications team to support this aim.</p> <p>Identify a lead officer to monitor internet and both local and social media.</p>	<p>20 July 2017</p> <p>20 July 2017</p> <p>Ongoing work</p>	<p>Lead Consultant Mental Health</p> <p>Local PH Mental Health Leads</p> <p>Local PH Mental Health Leads</p>

National Strategy				
		<p>Challenge stigma: Media campaign to support world suicide prevention day.</p> <p>Update council and stakeholder webpages to ensure effective signposting for those bereaved by suicide.</p>	<p>1 Sept 2017</p> <p>1 April 2017</p>	<p>Local PH Mental Health Leads</p> <p>Local PH Mental Health Leads</p>
6. Support research, data collection and monitoring		<p>Refresh Berkshire-wide suicide audit to include deaths during 2014-2016 to include data on GP consultations.</p> <p>To update data on the JSNA summary on suicide.</p>	<p>30 July 2017</p> <p>As per JSNA timetable</p>	<p>Local PH Mental Health Leads</p> <p>Local PH Mental Health Leads</p>

DIGITAL SOLUTIONS PROOF

References

All-Party Parliamentary Group on Suicide and Self-Harm Prevention. Inquiry into local suicide prevention plans in England. All-Party Parliamentary Group on Suicide and Self-Harm Prevention; 2015.

Barr, et al. Suicides associated with the 2008-10 economic recession in England: time trend analysis. *BMJ* 2012; 345:e5142 doi: 10.1136/bmj.e5142 (Published 14 August 2012).

Hawton K, Bergen H, Cooper J, Turnbull P, Waters K, Ness J, et al. Suicide following self-harm: findings from the multicentre study of self-harm in England: 2000-2012. *J Affect Disord.* 2015 Apr 1;175:147-51.

HM Government. Preventing suicide in England: A cross government strategy to save lives. London: Department of Health; 2012.

HM Government. Preventing suicide in England: Third progress report of the cross-government outcomes strategy to save lives. London: Department of Health; 2017.

Home Office. Multi-agency Statutory Guidance for the Conduct of Domestic Homicide Reviews. London: Home Office; 2016.

Making Mental Health Care Safer. The National Confidential Inquiry into Suicide and Homicide by People with Mental Illness. Annual Report and 20-year Review. University of Manchester; 2016. NHS England Mental Health Taskforce. The five year forward view for mental health. NHS England; 2016.

Office for National Statistics. Suicides in students <http://www.ons.gov.uk/ons/about-ons/what-we-do/publication-scheme/published-ad-hoc-data/health-and-social-care/november-2012/index.html>; 2012. [Accessed November 2016].

Office for National Statistics. Suicides in the UK in 2014. London: Office for National Statistics; 2015.

Office for National Statistics. Suicides in the UK in 2015. London: Office for National Statistics; 2016.

Public Health England (PHE). Local Suicide Prevention Planning, A Practical Resource. Public Health England; 2016.

Samaritans. Men, Suicide and Society: Why disadvantaged men in mid-life die by suicide. Ewell; Samaritans; 2012.

Sutherlands, R. PACTS: 26th Westminster Lecture for Samaritans on 'Working together to reduce suicide in transport'. 2015.

Appendix 1: Resources available

Factsheet on managing suicide risk in Primary Care http://www.connectingwithpeople.org/sites/default/files/SuicideMitigationInPrimaryCareFactsheet_0612.pdf

A free booklet on debt advice is available from:

<http://www.moneysavingexpert.com/credit-cards/mental-health-guide#collect>

Guide for health and social care workers to support people with debt and mental health problems written by the Royal College of Psychiatrists and Rethink Mental Illness:

<http://www.rcpsych.ac.uk/pdf/FinalDemandcf.pdf>

Primary Care Guidance on Debt and Mental Health from the Royal College of GPs and Royal College of Psychiatrists, due to be updated shortly: http://www.rcgp.org.uk/clinical/clinical-resources/~/_media/Files/CIRC/Mental%20health%20forum/Mental%20Health%20Page%20Sept%202013/PCMHF-Guidance-Debt-Mental-Health-Factsheet-2009.ashx

Leeds Bereavement Forum has produced a short document with details of local and national support services available. <http://www.leeds.gov.uk/docs/Bereavement%20leaflet%202013.pdf>

Grassroots Suicide Prevention Brighton & Hove Suicide Prevention Strategy Group provides an excellent website full of practical suicide prevention expertise.

http://prevent-suicide.org.uk/suicide_safer_brighton_and_hove.html

RAID service saves money as well as improving the health and well-being of its patients.

<http://www.bsmhft.nhs.uk/our-services/rapid-assessment-interface-and-discharge-raid/>

NHS Cornwall and Isles of Scilly, in partnership with Outlook South West

<http://www.outlooksw.co.uk/suicide-liaison-service>

Children and Young People's Mental Health Coalition Resilience and Results:

http://www.cypmhc.org.uk/resources/resilience_results/

State of Mind is a Rugby League mental health and wellbeing initiative which aims to raise awareness and tackle stigma. The organisation aims to reach men who may not normally contact health and social care services, and signpost them to where support is available. A round of Rugby League fixtures is dedicated to State of Mind, which maximises the publicity. The focus is on promoting player welfare and resilience in local communities. Super League players act as ambassadors reaching fans and amateur players through presentations, meetings and social networking, with positive messages being specially commissioned and tweeted. Films with specific themes are available at: www.stateofmindrugby.com

Samaritans Media Reporting Guidance:

<http://www.samaritans.org/media-centre/media-guidelines-reporting-suicide>

Appendix 2: Bracknell Forest Action Plan 2017-18

Areas for Action	Specific Risk Groups	Action in 2017-18	Timescale
Overarching Aims		Sign off / endorsement of this Berkshire Suicide Prevention Strategy by all Health & Wellbeing Boards in Berkshire.	1 April 2017
		All other local governance completed thus enabling this strategy to be owned by the Local Authorities and their partners.	1 April 2017
		Launch of strategy at multi-agency suicide prevention summit.	15 Oct. 2017
		Identify Suicide Prevention Champions to promote the Berkshire Strategy; and to work within the Local Authorities and partner agencies.	15 Oct. 2017
National Strategy			
1. Reduce the risk of suicide in key high-risk groups	Men	Promotion of CALM to a wider audience.	1 June 2017
	People in mental health care	Support BHFT in its Zero Suicide Approach.	Ongoing work
	Occupational Groups	Work with local Carers' support groups to highlight Mental Wellbeing issues and risk factors.	
	Carers (including young carers)	Multi agencies approach to identify individuals and sign posting for support/ local befriending service/ other services.	
	Socially isolated	Increase local befriender's awareness of Mental Wellbeing issues and Risk factors.	
2. Tailor approaches to improve mental health in specific groups	Community based approaches	Work with local Domestic Abuse Forum and Executive Group to provide support and information on suicide prevention.	
	People vulnerable due to economic circumstances	To share local Suicide Prevention strategy/ action plans/supporting materials with IAPT/ Job Centre and other employment support agencies. Increase agencies awareness of Mental Wellbeing issues and Risk factors.	

National Strategy			
3. Reduce access to the means of suicide		<p>Support Network Rail, British Transport police and The Samaritans with local escalation process locations and general suicide prevention work.</p> <p>Investigate suicides on council owned land and properties, and agree a local action plan.</p> <p>Analyse local data gathered from the suicide audit and/or real-time surveillance to identify trends and clusters and take appropriate action(s) e.g. work with local media.</p>	<p>Ongoing work</p> <p>15 Oct. 2017</p> <p>Ongoing work</p>
4. Provide better information and support to those bereaved or affected by suicide		<p>Promote resources for people bereaved and affected by suicide (e.g. Help is at Hand and National Suicide Prevention Alliance resources).</p>	<p>Ongoing work</p>
5. Support the media in delivering sensitive approaches to suicide and suicidal behaviour		<p>Convene a Berkshire-wide Summit on journalism and reporting standards with local press and media organisations, to develop and agree standards for reporting.</p> <p>Agree a local action plan with the local communications team to support this aim.</p> <p>Identify a lead officer to monitor internet and both local and social media.</p> <p>Challenge stigma: Media campaign to support world suicide prevention day.</p> <p>Update council and stakeholder webpages to ensure effective signposting for those bereaved by suicide.</p>	<p>20 July 2017</p> <p>20 July 2017</p> <p>Ongoing work</p> <p>10 Sept. 2017</p> <p>1 April 2017</p>
6. Support research, data collection and monitoring		<p>To update data on the JSNA summary on suicide.</p>	<p>As per JSNA timetable</p>

Appendix 3: Royal Borough of Windsor and Maidenhead Action Plan 2017-18

Areas for Action	Specific Risk Groups	Action in 2017-18	Timescale by:	Delivery Lead
Overarching Aims		Establish a multi-agency steering group: terms of reference to be agreed. Group will also be responsible for reviewing communication between primary and secondary care including risk assessment and escalation protocols.	Locally determined	To be locally determined
National Strategy				
1. Reduce the risk of suicide in key high-risk groups	Priority groups for 17/18: men; carers; unemployed; those who misuse substances; and those with mental health diagnoses.	<p>Build on existing local voluntary and community group programmes e.g. men in sheds.</p> <p>Training for gatekeepers relating to priority at-risk groups (Warwickshire).</p> <p>Support BHFT in its Zero Suicide Approach.</p> <p>Ensure adequate arrangements are in place for follow-up after discharge from secondary care.</p> <p>Consider strengths and issues arising from the Berkshire crisis concordat relating to the Royal Borough of Windsor and Maidenhead.</p>	<p>Ongoing work</p> <p>Locally determined</p> <p>Ongoing work</p> <p>Ongoing work</p> <p>Ongoing work</p>	
2. Tailor approaches to improve mental health in specific groups	Suicide prevention training	Map evidence of coverage by sector/organisation of self-harm and suicide prevention training.	Ongoing work	To be locally determined

National Strategy				
3.Reduce access to the means of suicide		<p>Support Network Rail, British Transport police and Samaritans with local escalation process locations and general suicide prevention work.</p> <p>Investigate suicides on council owned land and properties, and agree a local action plan.</p> <p>Analyse local data gathered from the suicide audit and/or real-time surveillance to identify trends and clusters and take appropriate action(s) e.g. work with local media.</p>	<p>Ongoing work</p> <p>15 Oct. 2017</p> <p>Ongoing work</p>	<p>To be locally determined</p> <p>To be locally determined</p> <p>To be locally determined</p>
4. Provide better information and support to those bereaved or affected by suicide		<p>Promote resources for people bereaved and affected by suicide (e.g. Help is at Hand and National Suicide Prevention Alliance resources).</p> <p>Update council and stakeholder webpages to ensure effective signposting for those bereaved by suicide.</p> <p>Map existing bereavement support and pathways.</p>	<p>Ongoing work</p> <p>Locally determined</p> <p>Ongoing work</p>	
5. Support the media in delivering sensitive approaches to suicide and suicidal behaviour		<p>Convene a Berkshire-wide Summit on journalism and reporting standards with local press and media organisations, to develop and agree standards for reporting.</p> <p>Agree a local action plan with the local communications team to support this aim.</p> <p>Identify a lead officer to monitor internet and both local and social media.</p> <p>Challenge stigma: Media campaign to support world suicide prevention day.</p> <p>Update council and stakeholder webpages to ensure effective signposting for those bereaved by suicide.</p>	<p>20 July 2017</p> <p>20 July 2017</p> <p>Ongoing work</p> <p>10 Sept. 2017</p> <p>1 April 2017</p>	

National Strategy				
6. Support research, data collection and monitoring		<p>To update data on the JSNA summary on suicide.</p> <p>Develop a suicide audit database (based on Bromley model) and continue to update relevant local data from sources which include: Office for National Statistics, Coroner's records, Thames Valley Police.</p> <p>Work with steering group members to review data about current levels of population need and service provision.</p> <p>Work with steering group members to map areas of high risk through information on locations of deaths and attempts. Take action to reduce suicide enablers (e.g. install signage, barriers) in line with evidence base.</p> <p>Undertake mapping relating to local suicide prevention and self-harm services.</p>	<p>As per JSNA timetable</p> <p>Locally determined</p> <p>Locally determined</p> <p>Locally determined</p> <p>Locally determined</p>	

DIGITAL SOLUTIONS PROOF

Appendix 4: Slough Action Plan 2017-18

Areas for Action	Specific Risk Groups	Action in 2017-18	Timescale by:
Overarching Aims		Sign off / endorsement of this Berkshire Suicide Prevention Strategy by all Health & Wellbeing Boards in Berkshire.	1 April 2017
		All other local governance completed thus enabling this strategy to be owned by the Local Authorities and their partners.	1 April 2017
		Launch of strategy at multi-agency suicide prevention summit.	15 Oct. 2017
		Identify Suicide Prevention Champions to promote the Berkshire Strategy; and to work within the Local Authorities and partner agencies.	15 Oct. 2017
National Strategy			
1. Reduce the risk of suicide in key high-risk groups	Men	Promotion of CALM to a wider audience.	Locally determined
	People who misuse substances	To partner with the drugs and alcohol team on reviewing the referral pathway for dual diagnosis. To continue to ensure that information on how to access DAAT services and seek help are readily available for young men.	
	People in mental health care	Support BHFT in its Zero Suicide Approach.	Ongoing work
	Occupational Groups	To support SME business on the Slough Trading Estate on incorporating mental health and wellbeing in their policies and advise on how to improve staff well-being; i.e.; promote resilience training.	

National Strategy			
2. Tailor approaches to improve mental health in specific groups	<p>Community based approaches</p> <p>Suicide prevention training</p> <p>People vulnerable due to economic circumstances</p> <p>Children and young people</p>	<p>To work with the community development team - to build community cohesion, etc.</p> <p>To identify and work with Housing and unemployment teams on MHFA training for staff.</p> <p>To deliver MHFA training to managers of SME businesses in Slough.</p> <p>To partner with NEET young people's team and train staff on MHFA.</p> <p>To design a service information leaflet for new migrant arrivals and to ensure that all frontline services have an access to the leaflet.</p> <p>To partner with young people service to design an intergenerational programme addressing loneliness and social isolation.</p>	
3. Reduce access to the means of suicide		<p>Support Network Rail, British Transport police and The Samaritans with local escalation process locations and general suicide prevention work.</p> <p>Investigate suicides on council owned land and properties, and agree a local action plan.</p> <p>Analyse local data gathered from the suicide audit and/or real-time surveillance to identify trends and clusters and take appropriate action(s) e.g. work with local media.</p>	<p>Ongoing work</p> <p>15 Oct. 2017</p> <p>Ongoing work</p>
4. Provide better information and support to those bereaved or affected by suicide		<p>Promote resources for people bereaved and affected by suicide (e.g. Help is at Hand and National Suicide Prevention Alliance resources).</p> <p>To conduct a mapping of services available for those that have been bereaved by suicide.</p> <p>Contact Samaritans SBCCG in order to identify Slough residents assessing the service and where they refer them to.</p>	<p>Ongoing work</p> <p>Locally determined</p>

National Strategy			
		To identify other local stakeholders and provide better information and support to those bereaved or affected by suicide.	
5. Support the media in delivering sensitive approaches to suicide and suicidal behaviour		<p>Convene a Berkshire-wide Summit on journalism and reporting standards with local press and media organisations, to develop and agree standards for reporting.</p> <p>Agree a local action plan with the local communications team to support this aim.</p> <p>Challenge stigma: Media campaign to support world suicide prevention day.</p> <p>Update council and stakeholder webpages to ensure effective signposting for those bereaved by suicide.</p>	<p>20 July 2017</p> <p>20 July 2017</p> <p>10 Sept. 2017</p>
6. Support research, data collection and monitoring		To update data on the JSNA summary on suicide.	As per JSNA timetable

DIGITAL SOLUTIONS PROOF

Appendix 5: Reading Action Plan 2017-18

What will be done - the task	Who will do it	By when	Outcome - the difference it will make	Supporting national indicators
Identify local sponsors to oversee Reading's Suicide Prevention Action Plan	Health & Wellbeing Board (Berkshire West Mental Health Strategy Group / Reading Mental Health Strategy Group)	February 2017	Reading actions to reduce deaths by suicide will be co-ordinated across agencies / There will be consistent local representation on the Berkshire Suicide Prevention Planning Group	
Develop a communication plan to raise awareness of Reading's Suicide Prevention Action Plan, including: - The formal launch of the Berkshire Suicide Prevention Strategy - Contributions to the 'Brighter Berkshire' Year of Mental Health 2017 - Marking World Suicide Prevention Day (10 September)	RBC Communications Team	April 2017	Individuals will have increased awareness of support available / Partners will know how to engage with and support the Reading Suicide Prevention Action Plan	
- Support the review of CALMzone and development of future commissioning plans for support services which target men - Review local DAAT contracts to ensure suicide prevention objectives are included - Develop post discharge support for people who have used mental health services via the Reading Recovery College	Wellbeing Team, RBC	October 2017 April 2017 Ongoing	Suicide risk will be mitigated for higher risk groups: men, people who abuse drugs or alcohol, people who have been in contact with mental health services	PHOF 4.10 - suicide rates

<p>Tailor approaches to improve mental health in specific groups:</p> <ul style="list-style-type: none"> - Support delivery of the local 'Future in Mind' programme to improve mental health in children and young people - Recognise the mental health needs of survivors and links to suicide prevention in the implementation of the Reading Domestic Abuse Strategy - Raise awareness of support available to survivors of sexual abuse through Trust House Reading - Contribute to a Berkshire wide review of targeted community based interventions, including suicide prevention and mental health first aid training 	<p>Local sponsors (see above)</p> <p>DENS, RBC</p> <p>Local sponsors (see above)</p> <p>Local sponsors (see above)</p>	<p>Ongoing</p> <p>tbc</p> <p>ongoing</p>	<p>Mental health will be improved for some specific groups (children and young people, survivors of domestic or sexual abuse) through tailored approaches.</p> <p>Future commissioning of community based interventions will be informed by a review of impact.</p>	
<p>- Analyse local data gathered from the suicide audit and/or real-time surveillance to identify trends and clusters and recommend appropriate action(s)</p>	<p>Wellbeing Team, RBC</p>	<p>ongoing</p>	<p>Access to the means of suicide will be reduced where possible.</p>	
<p>- Review pages on the Reading Services Guide to include national resources (e.g. 'Help is at Hand' and National Suicide Prevention Alliance resources) and signposting to local services</p> <p>- Map local bereavement support and access to specific support for bereavement through suicide</p>	<p>Wellbeing Team, RBC</p>	<p>June 2017</p>	<p>Those bereaved or affected by suicide will have access to better information and support.</p>	

<ul style="list-style-type: none"> - Ensure local media and communications staff are aware of Samaritans guidance on responsible suicide reporting - Support a Berkshire-wide Summit on journalism and reporting standards with local press and media organisations, to develop and agree standards for reporting. 	<p>Wellbeing Team, RBC</p>	<p>February 2017</p> <p>July 2017</p>	<p>Local media will be supported to report on suicide and suicidal behaviour in a sensitive manner.</p>	
<ul style="list-style-type: none"> - Update Reading JSNA module on suicide and self-harm - Refresh Reading Mental Health Needs Analysis 	<p>Wellbeing Team, RBC Adults Commissioning Team, RBC</p>	<p>tbc</p> <p>May 2016</p>	<p>Local and county-wide Suicide Prevention Action will be informed by up to date research, data collection and monitoring.</p>	

DIGITAL SOLUTIONS PROOF

Appendix 6: West Berkshire Action Plan 2017-18

Areas for Action	Specific Risk Groups	Action in 2017-18	Timescale
Overarching Aims		Sign off / endorsement of this Berkshire Suicide Prevention Strategy by all Health & Wellbeing Boards in Berkshire.	1 April 2017
		All other local governance completed thus enabling this strategy to be owned by the Local Authorities and their partners.	1 April 2017
		Launch of strategy at multi-agency suicide prevention summit.	15 Oct. 2017
		Identify Suicide Prevention Champions to promote the Berkshire Strategy; and to work within the Local Authorities and partner agencies.	15 Oct. 2017
		Set up local quarterly meetings to review the action plan.	Quarterly interval
National Strategy			
1. Reduce the risk of suicide in key high-risk groups	Men	Further development of "Pie and a pint" interventions.	Ongoing work
		Promotion of CALM to a wider audience.	Ongoing work
	People who self-harm	Monitor levels of self-harm.	
	People who misuse substances	Liaising with local substance misuse services.	
	People in mental health care	Support BHFT in its Zero Suicide Approach.	Ongoing work
2. Tailor approaches to improve mental health in specific groups	Community based approaches	Improve public awareness of suicide. Link with West Berkshire Emotional Health Academy.	
	Suicide prevention training	Delivery of Adult Mental Health First Aid Training.	

National Strategy			
	Children and young people	Delivery of Youth Mental Health First Aid Training and MHFA Schools Training	
3. Reduce access to the means of suicide		<p>Support Network Rail, British Transport police and The Samaritans with local escalation process locations and general suicide prevention work.</p> <p>Investigate suicides on council owned land and properties, and agree a local action plan.</p> <p>Analyse local data gathered from the suicide audit and/or real-time surveillance to identify trends and clusters and take appropriate action(s) e.g. work with local media.</p>	<p>Ongoing work</p> <p>15 October 2017</p> <p>Ongoing work</p>
4. Provide better information and support to those bereaved or affected by suicide		<p>Promote resources for people bereaved and affected by suicide (e.g. Help is at Hand and National Suicide Prevention Alliance resources)</p> <p>Seek views of those with lived experience on draft action plan</p> <p>Promotion of Newbury SOBs group</p>	<p>Ongoing work</p> <p>Locally determined</p> <p>Ongoing work</p>
5. Support the media in delivering sensitive approaches to suicide and suicidal behaviour		<p>Convene a Berkshire-wide Summit on journalism and reporting standards with local press and media organisations, to develop and agree standards for reporting.</p> <p>Agree a local action plan with the local communications team to support this aim.</p> <p>Identify a lead officer to monitor internet and both local and social media.</p> <p>Challenge stigma: Media campaign to support world suicide prevention day.</p> <p>Update council and stakeholder webpages to ensure effective signposting for those bereaved by suicide.</p>	<p>20 July 2017</p> <p>20 July 2017</p> <p>Ongoing work</p> <p>10 Sept. 2017</p> <p>1 April 2017</p>
6. Support research, data collection and monitoring		<p>To update data on the JSNA summary on suicide.</p> <p>Develop infographics to share with public.</p> <p>Link to W Berks mental health strategy.</p> <p>Link to W Berks health and wellbeing strategy.</p>	<p>As per JSNA timetable</p> <p>Locally determined</p>

Appendix 7: Wokingham Action Plan 2017-18

Areas for Action	Specific Risk Groups	Action in 2017-18	Timescale by:	Outcome Measure
Overarching Aims		Sign off / endorsement of this Berkshire Suicide Prevention Strategy by all Health & Wellbeing Boards in Berkshire.	1 April 2017	Signed copy of Strategy
		All other local governance completed thus enabling this strategy to be owned by the Local Authorities and their partners.	1 April 2017	Strategy implemented and agreed across the borough
		Launch of strategy at multi-agency suicide prevention summit.	15 Oct. 2017	High profile launch of strategy
		Identify Suicide Prevention Champions to promote the Berkshire Strategy; and to work within the Local Authorities and partner agencies.	15 Oct. 2017	Number of champions identified and trained across the partnership
National Strategy				
1. Reduce the risk of suicide in key high-risk groups	Men	Promotion of CALM to a wider audience.	1 June 2017	Widespread awareness of CALM and increase in numbers of men accessing the service
	People in mental health care	Support BHFT in its Zero Suicide Approach.	Ongoing work	Evidence of joint working and shared actions
	Occupational Groups	Awareness raising and training for local businesses on identifying early signs and how to respond.		Number of training sessions run
	LGBT groups	Working with local services such as TVPS.		Evidence of joint working and shared actions

	Carers (including young carers) and People with LTC	Work with local carer groups to raise awareness of Mental Health risks and prevention, promote local befriending and support groups.		Training provided. Information on readily available from carer groups and networks
	People who misuse substances	Work with the local treatment provider to ensure that risk of suicide and mental health are part of the assessment		Suicide risk and mental health area included in standard assessment
2. Tailor approaches to improve mental health in specific groups	Community based approaches	Engage with local groups such as faith groups and befriending services. Wellbeing work with tenants services.		Evidence of joint working and shared actions Evidence of joint working and shared actions. Information readily available to staff.
	Suicide prevention training	Plan and prioritise a programme of suicide prevention training and integrate into MECC work stream.		Training plan in place.
3. Reduce access to the means of suicide		Support Network Rail, British Transport police and Samaritans with local escalation process locations and general suicide prevention work.	Ongoing work	Robust prevention measures and escalation procedures are in place and all partners are aware of these.
		Investigate suicides on council owned land and properties, and agree a local action plan.	15 Oct. 2017	Case review process established and evidence of reports and actions taken.
		Analyse local data gathered from the suicide audit and/or real-time surveillance to identify trends and clusters and take appropriate action(s) e.g. work with local media.	Ongoing work	Data shared with partners.

<p>4. Provide better information and support to those bereaved or affected by suicide</p>		<p>Promote resources for people bereaved and affected by suicide (e.g. Help is at Hand and National Suicide Prevention Alliance resources).</p> <p>Review the availability of support for families and communities bereaved by suicide and affected by near misses.</p> <p>Promote the local Wokingham SOBS group, working with them to identify gaps.</p>	<p>Ongoing work</p> <p>Locally determined</p> <p>Ongoing work</p>	<p>Proportion of people referred to bereavement services</p> <p>Needs assessment carried out</p> <p>Evidence of promotional work and partnership working</p>
<p>5. Support the media in delivering sensitive approaches to suicide and suicidal behaviour</p>		<p>Convene a Berkshire-wide Summit on journalism and reporting standards with local press and media organisations, to develop and agree standards for reporting.</p> <p>Agree a local action plan with the local communications team to support this aim.</p> <p>Identify a lead officer to monitor internet and both local and social media.</p> <p>Challenge stigma: Media campaign to support world suicide prevention day.</p> <p>Update council and stakeholder webpages to ensure effective signposting for those bereaved by suicide.</p>	<p>20 July 2017</p> <p>20 July 2017</p> <p>Ongoing work</p> <p>1 Sept. 2017</p> <p>1 April 2017</p>	<p>Summit organised and reporting standards published. Reduced stigma around suicide and reduction in copycat suicides. Suicides are reported appropriately and sensitively.</p> <p>Communication Action Plan</p> <p>Officer identified</p> <p>Campaign held</p> <p>Webpages up to date and those bereaved access support</p>
<p>6. Support research, data collection and monitoring</p>		<p>To update data on the JSNA summary on suicide.</p>	<p>As per JSNA timetable</p>	<p>JSNA suicide chapter up to date</p>

Appendix 8: Membership of the Berkshire Suicide Prevention Steering Group as at December 2016

Angela Baker	Deputy Centre Director	PHE South East
Angus Tallini	GP and Mental Health lead for Berkshire West CCGs	Newbury & District NHS CCG
Anthony Barrett		NHS
Belinda Dixon		RBWM
Caroline Attard	Nurse Consultant, In-patient wards	Berkshire Healthcare Foundation NHS Trust
Carol-Anne Bidwell	Public Health Programme Officer	Wokingham Borough Council
Charlotte Ryall	Coroner's Officer	Reading Borough Council
Chris Allen		NHS
Colin Bibby		SEAP
Daren Bailey		Berkshire Healthcare Foundation NHS Trust
Darrell Gale	Consultant in Public Health	Wokingham Borough Council
Debbie Daly	Director of Nursing and Quality	NHS Berkshire West CCGs
Eugene Jones		Berkshire Healthcare Foundation NHS Trust
Geoff Dennis		Berkshire Healthcare Foundation NHS Trust
Gillian McGregor		Reading Council
Gwen Bonner	Clinical Director Reading Locality Clinical Director Research	Berkshire Healthcare Foundation NHS Trust
Helen Ranasinghe		Samaritans
Helena Fahie	Public Health Support Manager	PHE South East
Janette Searle	Preventative Services Development Manager	Reading Borough Council
Jason Jongali	Head of Mental Health & Learning Disability Commissioning	NHS Berkshire West CCGs
Jillian Hunt		Bracknell Forest Council
Jo Greengrass		NHS
Jonathan Groenen		Thames Valley Police
Julia Wales,		Slough Council
Kate Jahangard		Reading Council

Katie Simpson	GP and Mental Health lead for Berkshire East CCGs	NHS CCG
Ken Hikwa		Berkshire Healthcare Foundation NHS Trust
Kim McCall		Reading Borough Council
Lesley Wyman	Consultant in Public Health	West Berkshire Council
Lisa McNally	Consultant in Public Health	Bracknell Forest Council
Lise Llewellyn	Strategic Director of Public Health	Public Health Services Berkshire
Natalie Mears	Public Health Programme Officer	RBWM
Mark Spencer	Detective Chief Inspector; Deputy Commander - Slough	Thames Valley Police
Sally Murray	Head of Children's Commissioning	NHS Berkshire West CCGs
Nadia Barakat	Head of Mental Health & Learning Disabilities Commissioning	NHS Berkshire East CCGs
Nick Davies		RBWM
Rachel Johnson	Public Health Programme Officer	West Berkshire Council
Ramesh Kukar		Slough Council of Voluntary Services
Reva Stewart	Locality Director	Berkshire Healthcare Foundation NHS Trust
Richard Tredgett		Reading Samaritans
Rukayat Akanji-Suleman	Public Health Programme Officer	Slough Borough Council
Safron Simmonds	Project Manager	NHS Berkshire West CCGs
Sarah Bellars		NHS
Sue McLaughlin	Clinical Director / Nurse Consultant Slough locality	Berkshire Healthcare Foundation NHS Trust
Susanna Yeoman		Berkshire Healthcare Foundation NHS Trust
Tandra Forster	Head of Adult Social Care	West Berkshire Council
Tanya Demonne	Mental Health Coordinator, Safeguarding	Royal Berkshire Hospital Foundation NHS Trust
Timothy Foley		SEAP
Tony Dwyer		Berkshire Healthcare Foundation NHS Trust

DIGITAL SOLUTIONS PROOF



WOKINGHAM
BOROUGH COUNCIL



NHS
Newbury and District
Clinical Commissioning Group

NHS
North and West Reading
Clinical Commissioning Group

NHS
South Reading
Clinical Commissioning Group

NHS
Wokingham
Clinical Commissioning Group

DIGITAL SOLUTIONS PROOF

Address info here

